

c:\epodata\sca\eplogf\internal.log

1/1 - (C) WPI / DERWENT

AN - 2001-019312 [03]

AP - JP19990082660 19990326

CPY - TAIC

DC - B04 D13 D16 D21

FS - CFI

IC - A23K1/165 ; A23L1/30 ; A61K7/00 ; A61K38/46 ; A61P31/04 ; C12N9/36

MC - B04-B04M B04-D01 B04-L01 B14-A01 D03-H02 D03-M D05-A02 D08-A05 D08-B

M1 - [01] M423 M431 M710 M782 M905 P220 P912 Q233 Q254; RA00GC-T RA00GC-M  
RA00GC-N- [02] M423 M431 M710 M782 M905 P220 P912 Q233 Q254; RA0120-T RA0120-M  
RA0120-N

PA - (TAIC ) TAIYO KAGAKU KK

PN - JP2000270858 A 20001003 DW200103 C12N9/36 006pp

PR - JP19990082660 19990326

XA - C2001-005830

XIC - A23K-001/165 ; A23L-001/30 ; A61K-007/00 ; A61K-038/46 ; A61P-031/04 ;  
C12N-009/36AB - JP2000270858 NOVELTY - A lysozyme sugar composite (LSC) is isolated  
and purified from an egg white liquid containing at least one  
saccharide.

- ACTIVITY - Antimicrobial.

- Egg white liquid (EWL) (1 l) was homogenized, and EWL with protein content was isolated, to which trehalose (0.1 kg) was added and stirred for 1 hour at 5 deg. C to prepare the lysozyme sugar composite (LSC). EWL was treated with an alkali and then with a cation exchange resin like Amberlite IRC-50 (200 ml) to adsorb LSC. The resin settled separating a supernatant EWL was washed in water and 3% of salt solution (1000 ml) was added. 1N sodium hydroxide was added to elute LSC. The resin containing solution was filtered, and the obtained filtrate was set to pH 9.5. The salting method with 5% salt concentrations collected LSC (9.5 g) in a cake form was dried to obtain LSC crystals (3.1 g). Antimicrobial activity of LSC crystal opposing to gram negative microbe like Escherichia coli was determined by evaluating the number of living microbes for 24 hours at 35 deg. C. The result obtained showed that LSC crystal had 25% of the number of living microbes. Thus showing that LSC crystal had high antimicrobial activity opposing to E. coli.

- MECHANISM OF ACTION - None given.

- USE - As antimicrobial agent in gram positive and negative microbes, pharmaceutical, cosmetics, foodstuff and feed field, and as storage life improvement agent of foodstuffs such as boiled fish paste, custard cream, pickles, sausage, ham, cooked rice and egg processed goods. As quasi-drugs like mouth washing liquids and cosmetics like shampoo. Useful in livestock, preventing illness of aquatic animal feed additive, and animal drug. For inhibiting pathogenic microbe of dandruff and dental carries, and as rinse and dentifrice agent.

- ADVANTAGE - LSC effectively reinforce the antimicrobial activity opposing to the gram positive and negative microbes. The antimicrobial spectrum is effectively enlarged by LSC. LSC effectively reduces the density of the conventional anti-microbial compound. LSC as storage life improvement agent prevents decomposition and disinfection of

BEST AVAILABLE COPY

c:\cpodata\scaleplog\internal.log

foodstuffs. LSC as livestock prevents the illness caused by aquatic animal feed additive or animal drug.

- (Dwg. 0/0)

CN - RA00GC-T RA00GC-M RA00GC-N RA0120-T RA0120-M RA0120-N

IW - LYSOZYME SUGAR COMPOSITE ANTIMICROBIAL AGENT GRAM NEGATIVE MICROBE ISOLATE PURIFICATION EGG WHITE LIQUID CONTAIN

IKW - LYSOZYME SUGAR COMPOSITE ANTIMICROBIAL AGENT GRAM NEGATIVE MICROBE ISOLATE PURIFICATION EGG WHITE LIQUID CONTAIN

NC - 001

OPD - 1999-03-26

ORD - 2000-10-03

PAW - (TAIC ) TAIYO KAGAKU KK

TI - Lysozyme sugar composite as antimicrobial agent for gram negative microbe, is isolated and purified from egg white liquid containing saccharides

JP2000270858

## © WPI / DERWENT

- MC - B04-B04M B04-D01 B04-L01 B14-A01 D03-H02 D03-M D05-A02 D08-A05 D08-B
- AN - 2001-019312 [03]
- TI - Lysozyme sugar composite as antimicrobial agent for gram negative microbe, is isolated and purified from egg white liquid containing saccharides
- AB - JP2000270858 NOVELTY - A lysozyme sugar composite (LSC) is isolated and purified from an egg white liquid containing at least one saccharide.
- ACTIVITY Antimicrobial.
- Egg white liquid (EWL) (1 l) was homogenized, and EWL with protein content was isolated, to which trehalose (0.1 kg) was added and stirred for 1 hour at 5 deg. C to prepare the lysozyme sugar composite (LSC). EWL was treated with an alkali and then with a cation exchange resin like Amberlite IRC-50 (200 ml) to adsorb LSC. The resin settled separating a supernatant EWL was washed in water and 3% of salt solution (1000 ml) was added. 1N sodium hydroxide was added to elute LSC. The resin containing solution was filtered, and the obtained filtrate was set to pH 9.5. The salting method with 5% salt concentrations collected LSC (9.5 g) in a cake form was dried to obtain LSC crystals (3.1 g). Antimicrobial activity of LSC crystal opposing to gram negative microbe like Escherichia coli was determined by evaluating the number of living microbes for 24 hours at 35 deg. C. The result obtained showed that LSC crystal had 25% of the number of living microbes. Thus showing that LSC crystal had high antimicrobial activity opposing to E. coli.
- MECHANISM OF ACTION - None given.
- USE - As antimicrobial agent in gram positive and negative microbes, pharmaceutical, cosmetics, foodstuff and feed field, and as storage life improvement agent of foodstuffs such as boiled fish paste, custard cream, pickles, sausage, ham, cooked rice and egg processed goods. As quasi-drugs like mouth washing liquids and cosmetics like shampoo. Useful in livestock, preventing illness of aquatic animal feed additive, and animal drug. For inhibiting pathogenic microbe of dandruff and dental carries, and as rinse and dentifrice agent.
- ADVANTAGE - LSC effectively reinforce the antimicrobial activity opposing to the gram positive and negative microbes. The antimicrobial spectrum is effectively enlarged by LSC. LSC effectively reduces the density of the conventional antimicrobial compound. LSC as storage life improvement agent prevents decomposition and disintegration of foodstuffs. LSC as livestock prevents the illness caused by aquatic animal feed additive or animal drug.
- (Dwg. 0/0)
- IW - LYSOZYME SUGAR COMPOSITE ANTIMICROBIAL AGENT GRAM NEGATIVE MICROBE ISOLATE PURIFICATION EGG WHITE LIQUID CONTAIN
- PN - JP2000270858 A 20001003 DW200103 C12N9/36 006pp
- IC - A23K1/165 ;A23L1/30 ;A61K7/00 ;A61K38/46 ;A61P31/04 ;C12N9/36
- DC - B04 D13 D16 D21
- PA - (TAIC) TAIYO KAGAKU KK
- AP - JP19990082660 19990326
- JP - JP19990082660 19990326
- PD - 1999-03-26
- ORD - 2000-10-03

DEPT. OF AGRICULTURE